


FOR CERVICAL DYSTONIA AND  
CHRONIC SIALORRHEA IN ADULTS

## TAKING THE NEXT STEP WITH MYOBLOC®

Here's what you can expect and  
how to save on treatment



**MYOBLOC**®  
rimabotulinumtoxinB  
Injection [5,000 Units/mL]

Not actual patient.

### What is MYOBLOC?

MYOBLOC is a prescription medicine used in adults that is injected into:

- muscles and used to treat the abnormal head position and neck pain that happens with cervical dystonia (CD).
- glands that make saliva and is used to treat long-lasting (chronic) drooling (sialorrhea).

It is not known whether MYOBLOC is safe or effective in children.

### IMPORTANT SAFETY INFORMATION

#### What is the most important information I should know about MYOBLOC?

**MYOBLOC may cause serious side effects that can be life threatening. Call your doctor or get medical help right away if you have any of these problems after treatment with MYOBLOC:**

- **Problems swallowing, speaking, or breathing.** These problems can happen hours to weeks after an injection of MYOBLOC if the muscles that you use to breathe and swallow become weak and death can happen if you have severe problems.
  - People with breathing problems who use muscles in their neck to help them breathe may be at greater risk.
  - Swallowing problems may last several months and require a feeding tube. If severe, food or liquids may go into your lungs. People who already have swallowing or breathing problems are at highest risk.
- **Spread of toxin effects.** In some cases, botulinum toxin may affect areas of the body away from the injection site and cause symptoms of botulism which include: loss of strength and muscle weakness all over the body; trouble saying words clearly (dysarthria); double vision; loss of bladder control; blurred vision and drooping eyelids; trouble breathing; hoarseness or change or loss of voice (dysphonia); trouble swallowing (dysphagia).

**These symptoms can happen hours to weeks after you receive an injection of MYOBLOC.**

These problems could make it unsafe for you to drive a car or do other dangerous activities.

**Please see additional Important Safety Information throughout and full [Prescribing Information and Medication Guide](#), including Warning, and talk to your healthcare provider.**



## MYOBLOC<sup>®</sup> MAY BE ABLE TO HELP

MYOBLOC has been trusted by healthcare providers like yours for nearly a quarter century to treat cervical dystonia (CD) and chronic sialorrhea (CS) in adults, two conditions that can have a big impact on everyday life. **Whichever you're affected by, you and your doctor may decide to see if MYOBLOC can help.**

## WHAT TO EXPECT

MYOBLOC is an injection administered by your doctor in the office.



FOR CD,

MYOBLOC is injected into affected muscles allowing them to relax.



FOR CS,

MYOBLOC is injected into the salivary glands to block the release of a protein that increases saliva production.

After the first dose, another dose of MYOBLOC may be given after 12 weeks or longer, if it is needed. Your doctor may change the dose of MYOBLOC until the best dose is found.

### IMPORTANT SAFETY INFORMATION (cont'd)

#### Do not receive MYOBLOC if you:

- are allergic to MYOBLOC or any of the ingredients in MYOBLOC.
- had an allergic reaction to any other botulinum toxin product such as BOTOX<sup>®</sup>, BOTOX<sup>®</sup> Cosmetic (onabotulinumtoxinA), DYSPORT<sup>®</sup> (abobotulinumtoxinA), or XEOMIN<sup>®</sup> (incobotulinumtoxinA).
- have a skin infection at the planned injection site.

**Please see additional Important Safety Information on page 4 and full Prescribing Information and Medication Guide, including Warning, and talk to your healthcare provider.**

## HOW TO SAVE ON MYOBLOC® TREATMENT

If financial assistance in obtaining MYOBLOC treatment is needed, there are programs for eligible patients.



### MYOBLOC CO-PAY PROGRAM

With the MYOBLOC CO-PAY PROGRAM, eligible patients could receive up to \$4,000 each year to help with out-of-pocket expenses for MYOBLOC treatment.

#### ? WHO'S ELIGIBLE?

##### Patients who:

- Are 18 years or older and legal US residents
- Have commercial insurance coverage
- Are diagnosed with cervical dystonia (CD) or chronic sialorrhea (CS)
- Are NOT enrolled in a government insurance plan (eg, Medicare, Medicaid, TRICARE®, or other federal or state-funded programs)

#### ? WHAT'S COVERED?

- Any size vial of MYOBLOC
- Injection procedure costs
- Injection guidance costs –Ultrasound or EMG
- *Note: Residents of Michigan, Rhode Island, and Minnesota are not eligible for injection procedure or injection guidance costs*

#### ? HOW TO ENROLL

Speak to your healthcare provider. They will submit an enrollment form on your behalf in their online portal, or by mail or fax.

- No financial requirements

### FINANCIAL ASSISTANCE MAY BE AVAILABLE TO THOSE WHO QUALIFY

Talk to your healthcare provider to determine eligibility.

#### IMPORTANT SAFETY INFORMATION (cont'd)

**Before receiving MYOBLOC, tell your doctor about all your medical conditions, including if you have:**

a disease that affects your muscles and nerves (such as amyotrophic lateral sclerosis [ALS or Lou Gehrig's disease], myasthenia gravis or Lambert-Eaton syndrome); had any side effect from any botulinum toxin product in the past; a breathing problem, such as asthma or emphysema; a history of swallowing problems or inhaling food or fluid into your lungs (aspiration); bleeding problems; drooping eyelids; plans to have surgery; had surgery on your face.

**Please see additional Important Safety Information on page 4 and full [Prescribing Information](#) and [Medication Guide](#), including Warning, and talk to your healthcare provider.**



### What is MYOBLOC?

MYOBLOC® is a prescription medicine used in adults that is injected into:

- muscles and used to treat the abnormal head position and neck pain that happens with cervical dystonia (CD).
- glands that make saliva and is used to treat long-lasting (chronic) drooling (sialorrhea).

It is not known whether MYOBLOC is safe or effective in children.

### IMPORTANT SAFETY INFORMATION

#### What is the most important information I should know about MYOBLOC?

**MYOBLOC may cause serious side effects that can be life threatening. Call your doctor or get medical help right away if you have any of these problems after treatment with MYOBLOC:**

- **Problems swallowing, speaking, or breathing. These problems can happen hours to weeks after an injection of MYOBLOC** if the muscles that you use to breathe and swallow become weak and death can happen if you have severe problems.
  - People with breathing problems who use muscles in their neck to help them breathe may be at greater risk.
  - Swallowing problems may last several months and require a feeding tube. If severe, food or liquids may go into your lungs. People who already have swallowing or breathing problems are at highest risk.
- **Spread of toxin effects.** In some cases, botulinum toxin may affect areas of the body away from the injection site and cause symptoms of botulism which include: loss of strength and muscle weakness all over the body; trouble saying words clearly (dysarthria); double vision; loss of bladder control; blurred vision and drooping eyelids; trouble breathing; hoarseness or change or loss of voice (dysphonia); trouble swallowing (dysphagia).

**These symptoms can happen hours to weeks after you receive an injection of MYOBLOC.**

These problems could make it unsafe for you to drive a car or do other dangerous activities.

#### Do not receive MYOBLOC if you:

- are allergic to MYOBLOC or any of the ingredients in MYOBLOC.
- had an allergic reaction to any other botulinum toxin product such as BOTOX®, BOTOX® Cosmetic (onabotulinumtoxinA), DYSPORT® (abobotulinumtoxinA), or XEOMIN® (incobotulinumtoxinA).
- have a skin infection at the planned injection site.

**Before receiving MYOBLOC, tell your doctor about all your medical conditions, including if you have:** a disease that affects your muscles and nerves (such as amyotrophic lateral sclerosis [ALS or Lou Gehrig's disease], myasthenia gravis or Lambert-Eaton syndrome); had any side effect from any botulinum toxin product in the past; a breathing problem, such as asthma or emphysema; a history of swallowing problems or

inhaling food or fluid into your lungs (aspiration); bleeding problems; drooping eyelids; plans to have surgery; had surgery on your face.

#### Tell your doctor if you:

- are pregnant or plan to become pregnant. It is not known if MYOBLOC can harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if MYOBLOC passes into breast milk.

**Tell your doctor about all the medicines you take,** including prescription and over-the-counter medicines, vitamins and herbal supplements. **Talk to your doctor before you take any new medicine after you receive MYOBLOC.**

Using MYOBLOC with certain other medicines may cause serious side effects. **Do not start any new medicines until you have told your doctor that you have received MYOBLOC in the past. Especially tell your doctor if you:** have received any other botulinum toxin product in the last 4 months; have received injections of botulinum toxin such as BOTOX®, BOTOX® Cosmetic (onabotulinumtoxinA), DYSPORT® (abobotulinumtoxinA), or XEOMIN® (incobotulinumtoxinA) in the past. Be sure your doctor knows exactly which product you received; have recently received an antibiotic by injection; take muscle relaxants; take an allergy or cold medicine; take a sleep medicine.

#### What should I avoid while receiving MYOBLOC?

MYOBLOC may cause loss of strength or general muscle weakness or vision problems within hours to weeks of receiving MYOBLOC. **If this happens, do not drive a car, operate machinery or do other dangerous activities.**

#### What are the possible side effects of MYOBLOC?

**MYOBLOC can cause serious side effects including:**

- allergic reactions. Symptoms of an allergic reaction to MYOBLOC may include: itching, rash, redness, swelling, wheezing, trouble breathing, or dizziness or feeling faint. Tell your doctor or get medical help right away if you get wheezing, or trouble breathing, or if you get dizzy or faint.

**The most common side effects of MYOBLOC in people with cervical dystonia include** dry mouth; injection site discomfort or pain; trouble swallowing (dysphagia); headache.

**The most common side effects of MYOBLOC in people with sialorrhea include** dry mouth and trouble swallowing (dysphagia).

**You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088.**

**Please see full [Prescribing Information](#) and [Medication Guide](#), including Warning, and talk to your healthcare provider.**